

\*

Calculation of a function of complex variable by solving differential equations

Let us e.g. calculate  $\exp(z)$  from the differential equation  $\frac{du}{dt} = u$  with the boundary condition  $u(0) = 1$ . Introducing the real parameter  $s \in [0, 1]$  and making a substitution  $t = sz$ , we get  $u'(s) = zu(s)$ ,  $u(0) = 1$ , where the prime denotes  $\frac{d}{ds}$ . To avoid complex arithmetics we introduce  $v = \Re(u)$ ,  $w = \Im(u)$  and get a system of two real ordinary differential equations (with  $z = x + iy$ )

$$\begin{cases} v'(s) = xv(s) - yw(s), \\ w'(s) = xw(s) + yv(s), \\ v(0) = 1, \\ w(0) = 0, \end{cases}, \quad (1)$$

the solution to which gives the sought  $\exp(z)$ :  $\Re(\exp(z)) = v(1)$ ,  $\Im(\exp(z)) = w(1)$ .

This system can be solved with our routines `rkdriver/pcdriver`. For larger  $z$  one can first use

$$\exp(z_1 + z_2) = \exp(z_1) \exp(z_2), \quad (2)$$

$$\exp(n) = e^n, \quad \exp(i\pi/2) = i. \quad (3)$$

\*

Calculation of a function of complex variable by integration

Let us calculate e.g.  $\arctan(z)$  using the integral representation

$$\arctan(z) = \int_0^z \frac{dt}{1+t^2}. \quad (4)$$

The integration path can be taken as a straight line from 0 to  $z = x + iy$  as  $t = sz$ , where  $s \in [0, 1]$  is a real parameter,

$$\arctan(z) = z \int_0^1 \frac{ds}{1+(sz)^2}. \quad (5)$$

one can avoid using complex arithmetics by calculating separately the real and imaginary parts of the integral with our routine `adapt`. If  $|z| > 1$  one should perhaps instead use

$$\arctan(z) = \frac{\pi}{2} - \int_z^\infty \frac{dt}{1+t^2}. \quad (6)$$